



We create chemistry

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Opera[®]

Fungicide

**ACTIVE CONSTITUENT: 85 g/L PYRACLOSTROBIN
62.5 g/L EPOXICONAZOLE**

GROUP	3	11	FUNGICIDE
-------	----------	-----------	-----------

For the control of a broad range of fungal diseases in Wheat, Barley and Oat/Oaten hay as per the DIRECTIONS FOR USE table.

**IMPORTANT: READ THE SAFETY DIRECTIONS BEFORE
USING THIS PRODUCT**

CONTENTS: 5L, 10L, 20L 110L

BASF Australia Ltd

ABN 62 008 437 867

Level 12, 28 Freshwater Place Southbank VICTORIA 3006

crop-solutions.basf.com.au

® Registered trademark of BASF

APVMA Approval No.: 62753/115978

SE	FORMULATION TYPE Suspo- Emulsion
-----------	--

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply more than two applications of Opera (or any other strobilurin or Group 11 fungicide) in any one season on the same paddock.

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour at the application site.

DO NOT apply with spray droplets smaller than a MEDIUM spray droplet size category according to nozzle manufacturer specifications that refer to ASAE S572 standard or the BCPC guideline.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are: 1. Date with start and finish times of application; 2. Location address and paddock/s sprayed; 3. Full name of this product; 4. Amount of product used per hectare and number of hectares applied to; 5. Crop/situation and weed/pest; 6. Wind speed and direction during application; 7. Air temperature and humidity; 8. Nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; 9. Name and address of person applying this product. (Additional record details may be required by the State or territory where this product is used.)

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers downwind from the application area and within the mandatory no-spray zones shown in Table 1 below.

Table 1 – No-Spray Zones for Protection of the Aquatic Environment		
FOR AERIAL APPLICATION		
Wind Speed Range at Time of Application	Downwind Mandatory No-Spray Zone	
	Fixed-Wing	Helicopter
from 3 to 8 kilometres per hour	400 metres	250 metres
from 8 to 14 kilometres per hour	650 metres	350 metres
from 14 to 20 kilometres per hour	800 metres	450 metres
FOR GROUND APPLICATION		
from 3 to 20 kilometres per hour	40 metres	

DO NOT apply if there are livestock, pasture or any land that is producing feed for livestock downwind from the application area and within the mandatory no-spray zones shown in Table 2 below.

Table 2 – No-Spray Zones for Protection of International Trade	
FOR AERIAL APPLICATION	
Wind Speed Range at Time of Application	Downwind Mandatory No-Spray Zone
from 3 to 8 kilometres per hour	160 metres
from 9 to 20 kilometres per hour	180 metres
FOR GROUND APPLICATION	
from 3 to 20 kilometres per hour	10 metres

CROP	DISEASE	RATE/ha	CRITICAL COMMENTS
Wheat	Leaf rust (<i>Puccinia recondita</i>)	500-1000mL	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and completion of flag leaf emergence (Z32-41) if disease is present or conditions favour disease development. Apply at the higher rate when disease pressure is high. Apply a maximum of 1000mL/ha/crop when applying by air. DO NOT apply later than Z59 as yield responses are unlikely
	Stripe rust (<i>Puccinia striiformis</i>) Stem rust (<i>Puccinia graminis</i> f.sp. <i>tritici</i>) Septoria nodorum blotch (<i>Phaeosphaeria nodorum</i>)	500mL	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and completion of flag leaf emergence (Z32-41) if disease is present or conditions favour disease development. Repeat spraying may be required, particularly if infection occurs early. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT apply later than Z59 as yield responses are unlikely.
	Powdery mildew (<i>Blumeria graminis</i> f.sp. <i>tritici</i>)	500mL	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and completion of flag leaf emergence (Z32-41) if disease is present or conditions favour disease development. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT apply later than Z59 as yield responses are unlikely.
Barley	Leaf rust (<i>Puccinia hordei</i>)	500mL	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and completion of flag leaf emergence (Z32-41) if disease is present or conditions favour disease development. DO NOT apply later than Z59 as yield responses are unlikely.
	Spot form of net blotch (<i>Pyrenophora teres</i> f. sp. <i>maculata</i>) Net form of net blotch (<i>Pyrenophora teres</i> f. sp. <i>teres</i>)	500-1000mL	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and completion of flag leaf emergence (Z32-41) if disease is present or conditions favour disease development. Use the higher rate when disease pressure is high. Apply a maximum of 1000mL/ha/crop when applying by air. DO NOT apply later than Z59 as yield responses are unlikely.

CROP	DISEASE	RATE/ha	CRITICAL COMMENTS
	Leaf scald (<i>Rynchosporium secalis</i>)	500mL	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and completion of flag leaf emergence (Z32-41) if disease is present or conditions favour disease development. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT apply later than Z59 as yield responses are unlikely.
	Powdery mildew (<i>Blumeria graminis</i> f. sp. <i>hordei</i>)	500-1000mL	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and completion of flag leaf emergence (Z32-41) if disease is present or conditions favour disease development. Use the higher rate when disease pressure is high. Apply a maximum of 1000mL/ha/crop when applying by air. Repeat spraying may be required, particularly if the low rate is used and infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT apply later than Z59 as yield responses are unlikely.
Oats & Oaten Hay	Septoria leaf blotch (<i>Phaeosphaeria avenaria</i>)	500mL	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and completion of flag leaf emergence (Z32-41) if disease is present or conditions favour disease development. Repeat spraying may be required, particularly if infection occurs early. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT apply later than Z59 as yield responses are unlikely.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

GRAZING: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 WEEKS AFTER APPLICATION.

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

Livestock destined for export markets: The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

Export Slaughter Interval (ESI)

AFTER OBSERVING THE WITHHOLDING PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

GENERAL INSTRUCTIONS

Opera Fungicide is a broad spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results.

Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present as indicated in the Directions for Use.

MIXING

Opera Fungicide is a suspo-emulsion formulation. The addition of a non-ionic surfactant at recommended rates may assist in providing better coverage of sprayed surfaces. Add the product to the half filled spray tank while agitating. Continue to agitate while topping up the tank and during spraying.

APPLICATION

Ground Application

Apply in a water volume of between 80 and 100L/ha, using flat fan nozzles operating at around 50cm above the top of the crop. Use the higher water volume in crops with heavier canopies.

Aerial Application

Opera should be applied in a spray volume of 20 to 30L/ha and the aircraft calibrated to provide maximum coverage using MEDIUM droplet size to minimize off-target drift.

FUNGICIDE RESISTANCE WARNING

GROUP	3	11	FUNGICIDE
--------------	----------	-----------	------------------

OPERA is a member of the DMI and strobilurin groups of fungicides. For fungicide resistance management the product is a Group 3 and Group 11 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 and Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 2 or Group 11 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, BASF Australia accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

COMPATIBILITY

Opera Fungicide is a suspo-emulsion (SE) formulation. When using in a tank mix with other products, the following mix order should be observed;

1. Half fill the spray tank with water. Maintain constant agitation;

2. Add solid formulation types such as water dispersible granule (WDG/WG), wettable powder (WP), water soluble granule (SG) formulated products first and allow dispersion
3. Add simple liquid formulations such as soluble concentrate (SL) or liquid suspension formulations such as suspension concentrate (SC) formulations
4. Add any liquid formulations for dispersion such as emulsifiable concentrate (EC) and liquid formulation emulsions such as suspo-emulsion (SE) or micro-emulsion (ME) formulations
5. Add any water-soluble salts
6. Add any adjuvants as required
7. Add any foliar fertilizer as required
8. Add remaining water

For further information on compatibility please contact your local reseller or BASF representative.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NONTARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH CRUSTACEA AND ENVIRONMENT

Very toxic to aquatic life DO NOT contaminate streams, rivers, wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL (5L, 10L, 20L)

Store in the closed, original container in a dry, cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple, or preferably pressure, rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should NOT be burnt.

STORAGE AND DISPOSAL (110L only)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty container by pumping through the dry-break Micro Matic connection system. DO NOT attempt to unscrew the Micro Matic valve or breach the locked filling point. DO NOT contaminate the container with water or other foreign material. Ensure that the Micro Matic coupler, pump, meter and hoses are disconnected, triple rinsed with clean water and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. DO NOT dispose of undiluted chemicals on-site.



We create chemistry

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. May irritate the nose and throat. May irritate the skin. Will irritate the eyes. Repeated exposure may cause allergic disorders. Avoid inhaling vapour or spray mist. Avoid contact with eyes and skin. Repeated minor exposure may have a cumulative poisoning effect. When opening the container, mixing and loading and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and washable hat and elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS).

CONDITIONS OF SALE

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non-excludable conditions or warranties of the Competition and Consumer Act 2010 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

APVMA Approval No: 62753/115978

Batch No:

Date of Manufacture:

® = Registered trademark of BASF

© BASF 2023

BASF Australia Ltd

ABN 62 008 437 867

Level 12, 28 Freshwater Place

Southbank VICTORIA 3006

crop-solutions.basf.com.au

FOR SPECIALIST ADVICE IN AN

EMERGENCY ONLY

PHONE 1800 803 440

TOLL FREE - ALL HOURS - AUSTRALIA WIDE

Opera[®]

Fungicide

85 g/L PYRACLOSTROBIN
62.5 g/L EPOXICONAZOLE

Additional statements as required by Safe Work Australia in accordance with the Globally Harmonized System of Classification and Labelling (GHS)

HAZARD STATEMENT

Suspected of causing cancer. Suspected of damaging fertility. Suspected of damaging the unborn child. Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS (Prevention)

Use only outdoors or in a well-ventilated area. Do not handle until all safety precautions have been read and understood. Contaminated work clothing should not be allowed out of the workplace. Do not eat, drink or smoke when using this product.

PRECAUTIONARY STATEMENTS (Response)

IF exposed or concerned: Call a POISON CENTER or doctor/physician. IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF ON SKIN (or hair): Wash with plenty of soap and water. If skin irritation or rash occurs: Call a POISON CENTER or doctor/physician. IF SWALLOWED: rinse mouth. Collect spillage. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse.